

## ANTG ROCKY

### Consumer Medicine Information (CMI) summary

The full CMI on the next page has more details. If you are worried about using this medicine, speak to your doctor or pharmacist.

#### 1. WHY AM I USING ANTG ROCKY?

ANTG Rocky contains the active ingredient whole dried Cannabis sativa flower. ANTG Rocky is not approved for any specific indication and should only be used to treat conditions or symptoms advised by your doctor.

Your doctor might recommend its use to reduce nausea, vomiting, chronic non cancer pain, cancer related pain, spasticity from neurological conditions and muscle spasms, improve sleep and appetite, depression and anxiety. Your doctor might prescribe ANTG Rocky for another purpose.

For more information, see [Section 1. Why am I using ANTG Rocky?](#) in the full CMI.

#### 2. WHAT SHOULD I KNOW BEFORE I USE ANTG ROCKY?

Do not use if you have ever had an allergic reaction to cannabis or any of the ingredients listed at the end of the CMI.

**Talk to your doctor if you have any other medical conditions, take any other medicines, or are pregnant or plan to become pregnant or are breastfeeding.**

For more information, see [Section 2. What should I know before I use ANTG Rocky?](#) in the full CMI.

#### 3. WHAT IF I AM TAKING OTHER MEDICINES?

Some medicines may interfere with ANTG Rocky and affect how it works. ANTG Rocky may affect how other medicines work.

A list of these medicines is in [Section 3. What if I am taking other medicines?](#) in the full CMI.

#### 4. HOW DO I USE ANTG ROCKY?

Use the amount of ANTG Rocky directed by your doctor. Follow your doctor's instructions and use ANTG Rocky until your doctor tells you to stop.

More instructions can be found in [Section 4. How do I use ANTG Rocky?](#) in the full CMI.

## ANTG ROCKY

(/ˈrɒki/) **Active ingredient: whole dried Cannabis sativa flower**

### Consumer Medicine Information (CMI)

This leaflet provides important information about using ANTG Rocky. **You should also speak to your doctor or pharmacist if you would like further information or if you have any concerns or questions about using ANTG Rocky.**

#### WHERE TO FIND INFORMATION IN THIS LEAFLET:

1. [Why am I using ANTG Rocky?](#)
2. [What should I know before I use ANTG Rocky?](#)
3. [What if I am taking other medicines?](#)
4. [How do I use ANTG Rocky?](#)
5. [What should I know while using ANTG Rocky?](#)
6. [Are there any side effects?](#)
7. [Product details](#)

#### 1. WHY AM I USING ANTG ROCKY?

ANTG Rocky contains the active ingredient whole dried Cannabis sativa flower cured and trimmed from a hybrid cultivar of the cannabis sativa plant. The trichomes (glands) on the cannabis sativa flower are abundant with phytocannabinoid compounds. When taken internally, phytocannabinoids interact with the body's receptors to produce psychoactive effects (changes to a person's mental state, such as behavior, mood, thoughts or perception) and therapeutic effects. ANTG Rocky is not approved for any specific indication and should only be used to treat conditions or symptoms advised by your doctor.

Your doctor might recommend its use to reduce nausea, vomiting, chronic non cancer pain, cancer related pain, spasticity from neurological conditions and muscle spasms, improve sleep and appetite, and manage seizures, depression and anxiety. Your doctor might prescribe ANTG Rocky for another purpose. This medicine is only available with a doctor's prescription. Possession without authority is illegal.

#### 2. WHAT SHOULD I KNOW BEFORE I USE ANTG ROCKY?

##### WARNINGS

##### DO NOT USE ANTG ROCKY IF:

- You are allergic to any cannabis product or any of the ingredients listed at the end of this leaflet
- Always check the ingredients to make sure you can use this medicine
- You are less than 18 years old
- You have serious liver or kidney problems
- You have pre-existing heart problems, such as narrowing of the arteries (ischaemic or coronary heart disease), irregular heartbeat, poorly controlled high blood pressure, or have had previous heart attack or heart failure

#### 5. WHAT SHOULD I KNOW WHILE USING ANTG ROCKY?

<b>Things you should do</b>	<ul style="list-style-type: none"> <li>• Remind any doctor, dentist or pharmacist you visit that you are using ANTG Rocky.</li> <li>• You should use ANTG Rocky under medical supervision.</li> <li>• Talk to your healthcare provider if you do not feel better, or if you feel worse.</li> </ul>
<b>Things you should not do</b>	<ul style="list-style-type: none"> <li>• Do not change your dosage without first checking with your doctor.</li> </ul>
<b>Driving or using machines</b>	<ul style="list-style-type: none"> <li>• Do not drive or operate heavy or dangerous machines while being treated on this product.</li> </ul>
<b>Drinking alcohol</b>	<ul style="list-style-type: none"> <li>• Alcohol may increase the effects of ANTG Rocky and you should avoid taking it while taking this medicine. Alcohol taken in combination with ANTG Rocky can impair your cognitive abilities, coordination, reaction time and motor skills.</li> </ul>
<b>Looking after your medicine</b>	<ul style="list-style-type: none"> <li>• Store below 25°C and protect from light.</li> </ul>

For more information, see [Section 5. What should I know while using ANTG Rocky?](#) in the full CMI.

#### 6. ARE THERE ANY SIDE EFFECTS?

There are a number of side effects associated with this medicine. It is important to be aware of them so that you can identify any symptoms if they occur (see the full CMI for more details). Common, but usually mild side effects known from medicinal cannabis treatment (both cannabidiol (CBD) and delta-9-tetrahydrocannabinol (THC)) include fatigue and sedation, dizziness, nausea, dry mouth, and increase or decrease in appetite. Serious side effects of THC (and products high in THC) include convulsions, feeling 'high', depression, confusion, hallucinations, paranoid delusions, psychosis (severe mental condition where a person loses contact with reality, and is unable to think and judge clearly), cognitive distortion (having thoughts that are not true), and allergy symptoms.

For more information, including what to do if you have any side effects, see [Section 6. Are there any side effects?](#) in the full CMI.

#### CHECK WITH YOUR DOCTOR IF YOU:

- Have or have had any other medical conditions, especially the following:
  - Kidney problems
  - Liver problems
  - Serious mental disorders, such as schizophrenia or another psychotic disorder
  - Panic attacks
  - Lung problems
  - Heart problems
  - Epilepsy or recurrent seizures
- You have a history of dependence or addictive behaviour associated with prescription or non-prescription products
- You are 65 years or older

During treatment, you may be at risk of developing certain side effects. It is important you understand these risks and how to monitor for them. See additional information under [Section 6. Are there any side effects?](#)

#### PREGNANCY AND BREASTFEEDING

Check with your doctor if you are pregnant or intend to become pregnant.

Talk to your doctor if you are breastfeeding or intend to breastfeed. Cannabinoids can pass into breastmilk and may affect foetal development.

Do not take this product whilst breastfeeding until you and your doctor have discussed the risks and benefits involved.

#### USE IN CHILDREN

ANTG Rocky is not recommended for use in children or adolescents under 18 years of age.

#### MEDICAL SUPERVISION OF TREATMENT WITH ANTG ROCKY

ANTG Rocky treatment should be initiated and supervised by a qualified healthcare provider experienced in the treatment of the symptom or disorder for which it is prescribed.

#### 3. WHAT IF I AM TAKING OTHER MEDICINES?

Tell your doctor or pharmacist if you are taking any other medicines, including any medicines, vitamins or supplements that you buy without a prescription from your pharmacy, supermarket or health food shop.

**Some medicines may interfere with ANTG Rocky and affect how it works. ANTG Rocky may affect how other medicines work. Tell your doctor if you are using medicines for:**

- Cold or allergy (medicines that make you drowsy, such as antihistamines)
- Pain (medicines known as opioids, such as oxycodone, hydrocodone, codeine, and morphine)
- Muscle-related conditions, such as muscle spasms, overactive bladder, movement disorders (Parkinson's disease), and asthma (medicines known as anticholinergics, e.g. tolterodine, oxybutynin, darifenacin, solifenacin, benzotropine mesilate, trihexyphenidyl, ipratropium bromide, tiotropium bromide)
- Sleeping problems (sleeping pills or sedatives/hypnotics, e.g. benzodiazepine, zopiclone, zolpidem)
- Epilepsy, seizures, or nerve pain (e.g. carbamazepine, phenytoin, and phenobarbital)
- Attention Deficit Hyperactivity Disorder (ADHD) or narcolepsy (medicines such as amphetamines that may increase your heart rate and blood pressure)

- Bacterial or fungal infections (e.g. rifampicin, clarithromycin, ketoconazole, itraconazole, amphotericin B)
- Suppression of the immune system (e.g. cyclosporine)
- Thinning of your blood (medicines known as coumarins, e.g. warfarin)

**Check with your doctor or pharmacist if you are not sure about what medicines, vitamins or supplements you are taking and if these affect ANTG Rocky.**

#### 4. HOW DO I USE ANTG ROCKY?

##### HOW MUCH TO USE

- Use the amount of ANTG Rocky directed by your doctor.
- The amount of ANTG Rocky your doctor tells you to use each day will depend on your condition, whether you are taking any other medicines, as well as how you respond to and tolerate the product.
- Your doctor may recommend that you start with a low dose of ANTG Rocky and then slowly increase the dose to the lowest amount needed to improve your symptoms.
- Follow the instructions provided and use ANTG Rocky until your doctor tells you to stop.

##### WHEN TO TAKE ANTG ROCKY

- ANTG Rocky can be taken with or without food.
- Taking ANTG Rocky under similar conditions (e.g. before or after food/meals) each day may help achieve a more consistent effect.

##### HOW TO TAKE ANTG ROCKY

- This product should be administered as directed by your doctor.
- Effects from ANTG Rocky may vary depending on the dose taken, how long you have been taking this medicine, the amount and type of food you have had when taking this medicine, and your individual response.

##### IF YOU FORGET TO USE ANTG ROCKY

ANTG Rocky should be used regularly at the same time each day. If you miss your dose at the usual time, take it as soon as you remember, then go back to taking this product as you would normally.

**If you have missed a dose by more than 4 hours, skip the dose you missed and take your next dose when you are meant to.**

**Do not take a double dose to make up for the dose you missed.**

##### IF YOU USE TOO MUCH ANTG ROCKY

If you think that you have used too much ANTG Rocky, you may need urgent medical attention.

##### You should immediately:

- phone the National Poisons Centre (**by calling 0800 764 766**), or
- contact your doctor, or
- go to the Emergency Department at your nearest hospital.

**You should do this even if there are no signs of discomfort or poisoning.**

#### 5. WHAT SHOULD I KNOW WHILE USING ANTG ROCKY?

##### THINGS YOU SHOULD DO

- You should use ANTG Rocky under medical supervision to help determine if the product is working or if you are having any side effects from the treatment.
- Talk to your healthcare provider if you do not feel better, or if you feel worse

Remind any doctor, dentist or pharmacist you visit that you are using ANTG Rocky.

##### THINGS YOU SHOULD NOT DO

- Do not change your dosage without first checking with your doctor.

##### TRAVELLING WITH ANTG ROCKY

- Before travelling to another country (this includes your final destination and any in transit countries), consult with your doctor, country consulates, The Ministry of Foreign Affairs and Trade, insurance company and/or pharmacist to confirm if it is legal for you to travel with this product.

##### DRIVING OR USING MACHINES

**Do not drive or operate heavy or dangerous machines while being treated on this product. Talk to your doctor on when to resume these activities after completing treatment.**

##### DRINKING ALCOHOL

**Tell your doctor if you drink alcohol.**

Alcohol may increase the effects of ANTG Rocky and you should avoid taking it while taking this medicine. Alcohol taken in combination with ANTG Rocky can impair your cognitive abilities, coordination, reaction time and motor skills.

#### LOOKING AFTER YOUR MEDICINE

- Store below 25°C.
- Follow the instructions in the carton on how to take care of your medicine properly.
- Store it in a cool dry place away from moisture, heat or sunlight; for example, do not store it:
  - in the bathroom or near a sink, or
  - in the car or on window sills.

**Keep it where young children cannot reach it.**

##### GETTING RID OF ANY UNWANTED MEDICINE

If you no longer need to use this medicine or it is out of date, take it to any pharmacy for safe disposal.

Do not use this medicine after the expiry date.

#### 6. ARE THERE ANY SIDE EFFECTS?

All medicines can have side effects. If you do experience any side effects, most of them are minor and temporary. However, some side effects may need medical attention.

See the information below and, if you need to, ask your doctor or pharmacist if you have any further questions about side effects.

#### SIDE EFFECTS

Side effects	What to do
<b>Changes to mental status:</b> <ul style="list-style-type: none"> <li>• Poor short-term memory</li> <li>• Lack of concentration</li> <li>• Confusion</li> <li>• Abnormal thoughts</li> <li>• Poor coordination and balance</li> <li>• Dizziness</li> </ul>	Speak to your doctor if you have any of these less serious side effects and they worry you.
<b>Personality or behavioural changes:</b> <ul style="list-style-type: none"> <li>• Anxiety</li> <li>• Loss of inhibition</li> <li>• Quiet and reflective mood, or other unusual changes in mood</li> <li>• Changes in appetite</li> <li>• Spontaneous laughter</li> <li>• Sleepiness/feeling tired</li> </ul>	
<b>Eye-related:</b> <ul style="list-style-type: none"> <li>• Bloodshot eyes</li> <li>• Altered visual perception</li> </ul>	
<b>Ear-related:</b> <ul style="list-style-type: none"> <li>• Altered hearing perception</li> </ul>	
<b>General body-related:</b> <ul style="list-style-type: none"> <li>• Phlegm and cough</li> <li>• Dry mouth</li> <li>• Malaise (feeling unwell) or a general feeling of discomfort</li> <li>• Nausea</li> <li>• Muscle weakness</li> </ul>	Speak to your doctor if you have any of these less serious side effects and they worry you.

#### SERIOUS SIDE EFFECTS

Serious side effects	What to do
<b>Changes to mental status or personality:</b> <ul style="list-style-type: none"> <li>• Depression or thoughts of suicide</li> <li>• Paranoia</li> <li>• Seeing or hearing things that are not there</li> <li>• Irrational thinking</li> <li>• Hallucinations</li> <li>• Addiction/dependence</li> </ul>	Call your doctor straight away, or go straight to the Emergency Department at your nearest hospital if you notice any of these serious side effects.
<b>Heart-related:</b> <ul style="list-style-type: none"> <li>• Rapid or irregular heartbeat</li> <li>• Signs of low blood pressure, such as light-headedness, dizziness or fainting</li> </ul>	
<b>Nervous system-related:</b> <ul style="list-style-type: none"> <li>• Seizures or convulsions (if you do not have epilepsy)</li> </ul>	
<b>Allergic reaction-related:</b> <ul style="list-style-type: none"> <li>• Shortness of breath</li> <li>• Wheezing or difficulty breathing</li> <li>• Swelling of the face, lips, tongue or other parts of the body</li> <li>• Rash, hives or swelling on the skin</li> </ul>	

**Tell your doctor or pharmacist if you notice anything else that may be making you feel unwell.**

Other side effects not listed here may occur in some people.

#### REPORTING SIDE EFFECTS

After you have received medical advice for any side effects you experience, you can report side effects to your doctor, pharmacist and/or NUBU Pharmaceuticals. By reporting side effects, you can help provide more information on the safety of this medicine.

**Always make sure you speak to your doctor or pharmacist before you decide to stop taking any of your medicines.**

#### 7. PRODUCT DETAILS

This medicine is only available with a doctor's prescription.

##### WHAT ANTG ROCKY CONTAINS

<b>Active ingredient (main ingredient)</b>	Each bottle has 10 g whole dried Cannabis sativa flower, which contains 2500mg of delta-9- tetrahydrocannabinol (25% THC), cannabidiol (<1% CBD) and other minor cannabinoids.
<b>Other ingredients (inactive ingredients)</b>	N/A

Do not take this medicine if you are allergic to any of these ingredients.

##### WHAT ANTG ROCKY LOOKS LIKE

Approximately 1–5 cm, brown-green clustered flowers with a characteristic cannabis smell. Available in a white HDPE jar with a child-resistant cap and tamper proof seal.

##### WHO DISTRIBUTES ANTG ROCKY

NUBU Pharmaceuticals

Auckland, New Zealand.

Byron Bay NSW 2481 Australia

Email: enquiries@nubupharma.com

Ph: 0800 46 33 22

Date of Preparation January 2022. NUBU024V1.1